

Classifying the irreducible components of moduli stacks of torsion-free sheaves on K3 surfaces and an application to Brill-Noether theory

Yuki Mizuno
(水野 雄貴)

Waseda University

October 21, 2020

Introduction

Purpose

Irr decomp of **moduli stacks of torsion-free sheaves**
of rk 2 on K3 surfaces of $\rho = 1$

&

irr decomp of **Brill-Noether(BN) locus** on Hilbert schs of pts.

※ Moduli stacks can parametrize **unstable sheaves**.

Previous research

- **The case of ruled surfaces**
→ C.Walter (1995)
- **Stratification of moduli stacks**
→ V.Hoskins (2018) or T.L.Gomez, I.Sols and A.Zamora (2015)

Mukai vector

X : Proj K3 surf/ \mathbb{C} of $\rho = 1$, $E \in \text{Coh}(X)$

1. $\mathbf{v}(E) := (\text{rk}(E), c_1(E), \text{ch}_2(E) + \text{rk}(E)) \in \mathbb{Z} \oplus \text{Pic}(X) \oplus \mathbb{Z}$
2. $\langle \mathbf{v}, \mathbf{w} \rangle := -[\mathbf{v}]_0[\mathbf{w}]_2 + [\mathbf{v}]_1[\mathbf{w}]_1 - [\mathbf{v}]_2[\mathbf{w}]_0 \in \mathbb{Z}$
, where $\mathbf{v} := ([\mathbf{v}]_0, [\mathbf{v}]_1, [\mathbf{v}]_2) \in \mathbb{Z} \oplus \text{Pic}(X) \oplus \mathbb{Z}$

Moduli stacks

3. $\mathcal{M}(v)$:

Ob. flat family \mathcal{E}/U paramet torsion-free sheaves w/
Mukai vector v

Mor. $(\varphi, \alpha) : \mathcal{E}/U \rightarrow \mathcal{E}'/U'$
 $(\varphi : U \rightarrow U' : \text{mor of schs}, \alpha : \mathcal{E} \rightarrow (\text{id}_X \times \varphi)^* \mathcal{E}' : \text{iso})$

4. $\mathcal{M}_{(v_1, v_2)}^{\text{HN}}(v) := \left\{ E \in \mathcal{M}(v) \mid \begin{array}{l} \exists (0 \subset E_1 \subset E) : \text{HN-filtration} \\ \text{s.t. } v(E_1) = v_1, v(E/E_1) = v_2 \end{array} \right\}$

5. $\mathcal{M}^{\text{ss}}(v) := \{E \in \mathcal{M}(v) \mid E : \text{semistable}\}$

Irreducible decomposition of $\mathcal{M}(v)$

Main Theorem 1

v_0 : primitive Mukai vector

$v := mv_0$ ($m \in \mathbb{Z}$)

Assume $[v]_0 = 2$ & v satisfies one of (a) \sim (c)

- (a) : $\langle v, v \rangle > 0$, (b) : $\langle v, v \rangle = 0, -2$ and v is primitive ,
(c) : $\langle v, v \rangle < -2$ and $\langle v_0, v_0 \rangle \neq -2$.

Then,

$$\mathcal{M}(v) = \begin{cases} \overline{\mathcal{M}^{\text{ss}}(v)} \cup \bigcup_{\langle v_1, v_2 \rangle \leq 1} \overline{\mathcal{M}_{(v_1, v_2)}^{\text{HN}}(v)} & \text{(a), (b)} \\ \bigcup \overline{\mathcal{M}_{(v_1, v_2)}^{\text{HN}}(v)} & \text{(c)} \end{cases}$$

Irreducible decomposition of $\mathcal{M}(v)$

Remark

For proof of Thm 1,

- theory of stratification via HN-filt
- theory of moduli sps of sheaves on K3 surfs by K.Yoshioka

are important.

\rightsquigarrow We get the relation between the stratas by calculating dims etc..

Application to BN theory

Definition (BN locus of Hilbert schs of pts)

D : eff div on X

$N \in \mathbb{N}$ s.t. $N \leq h^0(\mathcal{O}(D))$

$$W_N^i(D) := \{Z \in \text{Hilb}^N(X) \mid h^1(\mathcal{I}_Z(D)) \geq i + 1\}$$

Remark

$H^0(\mathcal{I}_Z(D)) - \{0\}/\mathbb{C}^* = \text{eff divs lin equiv to } D \text{ passing through } Z.$

For general $Z \in \text{Hilb}^N(X)$,

$$h^0(\mathcal{I}_Z(D)) = h^0(\mathcal{O}_X(D)) - \ell(\mathcal{O}_Z) = \text{expected dimension.}$$

But, for $Z \in W_N^i(D)$,

$$h^0(\mathcal{I}_Z(D)) > h^0(\mathcal{O}_X(D)) - \ell(\mathcal{O}_Z).$$

$$\ast \quad h^0(\mathcal{I}_Z(D)) = h^0(\mathcal{O}_X(D)) - \ell(\mathcal{O}_Z) + h^1(\mathcal{I}_Z(D)).$$

Application to BN theory

Main Theorem 2

$D := nH$, $\mathbf{v} := (2, nH, \frac{n^2 H^2}{2} - N + 2)$, where H : amp gen of $\text{Pic}(X)$.
If $\langle \mathbf{v}, \mathbf{v} \rangle > 0$, there exists the following 1 to 1 corresp

{ the irr comps of $W_N^0(D)$ }

1 to 1 \updownarrow

$\{ \overline{\mathcal{M}_{(\mathbf{v}_1, \mathbf{v}_2)}^{\text{HN}}(\mathbf{v})} \mid (\mathbf{v}_1, \mathbf{v}_2) \text{ satisfying } (*) \} \cup \{ \overline{\mathcal{M}^{\text{ss}}(\mathbf{v})} \}$
 $\subsetneq \{ \text{the irr comps of } \mathcal{M}(\mathbf{v}) \}$.

$[v_1]_1, [v_2]_1 \neq 0$: effective, $\langle \mathbf{v}_1, \mathbf{v}_2 \rangle \leq 1$, $[v_2]_2 \geq -1$ (*)

Remark

- If Z : general in $W_N^0(D)$, the corresp is given by ext'n

$$0 \rightarrow \mathcal{O}_X \rightarrow E \rightarrow \mathcal{I}_Z(D) \rightarrow 0.$$

- Thm 2 \rightsquigarrow (non-emptiness,) the dims and the num of irr comps of $W_N^0(D)$.

Thank you for listening !